

TIPS FROM OUR READERS

A safe holder for adjusting indirect prostheses:
The silicone cube technique



Kelvin I. Afrashtehfar, DDS, MSc,^a Daniel Buser, DMD, Prof Dr med dent,^b and
Urs C. Belser, DMD, Prof Dr med dent^c



Figure 1. Patient with provisionally cemented maxillary 3-unit ceramic (zirconia framework, layered veneering) FDP (right central to left lateral incisor) before and after implant placement in the pontic area. A, Initial condition. B, Silicone holder. C, Pontic area and prosthetic margins to be reduced. D, Modified prosthesis replaced temporarily for healing phase.

^aImplant Fellow, Department of Reconstructive Dentistry and Gerodontology, School of Dental Medicine, University of Bern, Bern, Switzerland; and Visiting Research Associate, Department of Oral Surgery and Stomatology, School of Dental Medicine, University of Bern, Bern, Switzerland.

^bProfessor and Chairman, Department of Oral Surgery and Stomatology, School of Dental Medicine, University of Bern, Bern, Switzerland.

^cChairman Emeritus, Division of Fixed Prosthodontics and Biomaterials, School of Dental Medicine, University of Geneva, Geneva; and Guest Professor, Department of Reconstructive Dentistry and Gerodontology, School of Dental Medicine, University of Bern, Bern, Switzerland.




Figure 2. Two implant-retained maxillary interim prosthesis. A, Buccal surface of PMMA (polymethylmetacrylate) interim crown on maxillary left lateral incisor requires polishing. B, Silicone holder. C, Buccal aspect exposed for finishing and polishing. D, Polished interim prosthesis.

Small ceramic and interim restorations may be difficult to hold to adjust or polish, especially when wet. This technique uses polyvinyl siloxane to hold dental restorations securely during extraoral adjustments with rotary instruments.

A temporarily cemented maxillary 3-unit fixed partial denture with a zirconia framework and layered porcelain (Fig. 1A) required removal and adjustment in the course of implant placement in the pontic site. After fabrication of the silicone cube, the pontic area was reduced, and the margins were adjusted after coronally repositioned flap procedures. The modified prosthesis was adjusted¹⁻⁷ and temporarily seated immediately after the planned surgical procedures.

The second example is a maxillary PMMA (polymethylmetacrylate) interim prosthesis that lacked a well-polished facial surface (Fig. 2A). A silicone cube holder was fabricated for finishing and polishing this area with pumice and diamond polishing paste.⁸

PROCEDURE

1. Mix the polyvinyl siloxane (Sil-Tech Super; Ivoclar Vivadent AG, President Putty; Coltène) homogeneously according to the manufacturer's instructions; form a cube; and embed the restoration (Fig. 1B).

2. Ensure that the silicone material wraps around the restoration and exposes only the surface to be adjusted.
3. When the material has polymerized, cut and trim the polyvinyl siloxane excess with a scalpel (carbon steel scalpel blade #15; B. Braun Melsungen AG).
4. Upon completion of the adjustments (Figs. 1C, 2C), use the scalpel to gently remove the restoration from the silicone cube if necessary (Figs. 1D, 2D).

REFERENCES

1. Işeri U, Ozkurt Z, Yalnız A, Kazazoğlu E. Comparison of different grinding procedures on the flexural strength of zirconia. *J Prosthet Dent* 2012;107:309-15.
2. Hatanaka GR, Polli GS, Fais LM, Reis JM, Pinelli LA. Zirconia changes after grinding and regeneration firing. *J Prosthet Dent* 2017;118:61-8.
3. Bartolo D, Cassar G, Al-Haj Husain N, Özcan M, Camilleri J. Effect of polishing procedures and hydrothermal aging on wear characteristics and phase transformation of zirconium dioxide. *J Prosthet Dent* 2017;117:545-51.
4. Mundhe K, Jain V, Pruthi G, Shah N. Clinical study to evaluate the wear of natural enamel antagonist to zirconia and metal ceramic crowns. *J Prosthet Dent* 2015;114:358-63.
5. Mohammadi-Bassir M, Jamshidian M, Rezvani MB, Babasafari M. Effect of coarse grinding, overglazing, and 2 polishing systems on the flexural strength, surface roughness, and phase transformation of yttrium-stabilized tetragonal zirconia. *J Prosthet Dent* 2017;118:658-65.
6. Park C, Vang MS, Park SW, Lim HP. Effect of various polishing systems on the surface roughness and phase transformation of zirconia and the durability of the polishing systems. *J Prosthet Dent* 2017;117:430-7.
7. Clayton JA, Green E. Roughness of pontic materials and dental plaque. *J Prosthet Dent* 1970;23:407-11.

8. Guler AU, Kurt S, Kulunk T. Effects of various finishing procedures on the staining of provisional restorative materials. *J Prosthet Dent* 2005;93:453-8.

Corresponding author:

Dr Kelvin Ian Afrashtehfar
Department of Reconstructive Dentistry and Gerodontology
School of Dental Medicine, Faculty of Medicine
University of Bern
Rm C305, Freiburgstrasse 7, 3010 Berne
SWITZERLAND
Email: kelvin.afrashtehfar@zmk.unibe.ch

Acknowledgments

Dr Afrashtehfar thanks Prof Dr UC Belser, Guest Professor, Department of Reconstructive Dentistry and Gerodontology and Prof Dr D Buser, Chairman of the Department of Oral Surgery and Stomatology, ZMK Bern, for their mentorship during the International Team for Implantology Scholar year; and Prof Dr U Braegger, Chairman, Department of Reconstructive Dentistry and Gerodontology, ZMK Bern, and the International Team for Implantology, Basel, Switzerland, for their educational support.

Copyright © 2017 by the Editorial Council for *The Journal of Prosthetic Dentistry*.

Noteworthy Abstracts of the Current Literature

Analysis of endodontic complications following fixed prosthodontic rehabilitation

Uzgur Z, Uzgur R, Çolak H, Ercan E, Dalli M
Int J Prosthodont 2016; 29:565-9

Purpose. The aim of this study was to determine endodontic treatment needs and types of endodontic disease following fixed prosthodontic treatment 24 hours after tooth preparation, 1 week after tooth preparation, 1 month after placement, and 6 months after placement.

Material and methods. Study groups consisted of patients who attended a university dental hospital department of prosthodontics for fixed prosthodontic treatment from January 2011 to December 2013. All teeth were clinically and radiographically evaluated according to American Association of Endodontists evaluation criteria before preparation. Metal-ceramic fixed partial dentures were placed for all patients. A total of 1,633 abutment teeth were prepared with 1,100 pontics in 524 patients (214 female and 310 male). Participant age, sex, and tooth number were recorded. Endodontic treatment follow-up was scheduled for 24 hours after tooth preparation, 1 week after preparation, 1 month after placement, and 6 months after placement, and all teeth were evaluated after placement of FPDs according to a modified criteria.

Results. 2,733 retainers were placed with 624 FPDs. Of the FPDs, 332 (53%) were placed in the posterior and 196 (31.5%) in the anterior region. The remaining 96 FPDs (15.5%) were placed anteroposterior. The abutment/pontic ratio was 1.44:1. The number of retainers per FPD was 4.37. Of 1,633 abutment teeth, 103 were endodontically treated after placement of FPDs. Most observed endodontic disease was symptomatic irreversible pulpitis. There were statistically significant differences in terms of teeth regions ($P < .001$). When follow-up times of 24 hours, 1 week, 1 month, and 6 months were evaluated, there was no statistically significant difference among all teeth groups ($P > .05$).

Conclusions. A total of 2,733 retainers on 624 FPDs were evaluated over 6 months, and the mean endodontic treatment need ratio was 6.3%.

Reprinted with permission of Quintessence Publishing.