


Samir Haj Bloukh and Zehra Edis\*

# Crystal structure and antimicrobial properties of (1,4,7,10-tetraoxacyclododecane- $\kappa^4 O, O', O'', O'''$ )cesium(I) penta-iodide, $C_{16}H_{32}CsI_5O_8$



the atoms including atomic coordinates and displacement parameters.

**Table 1:** Data collection and handling.

Crystal:	Black cuboid
Size:	0.30 × 0.30 × 0.10 mm
Wavelength:	Mo K $\alpha$ radiation (0.71073 Å)
$\mu$ :	7.33 mm <sup>-1</sup>
Diffractometer, scan mode:	PDS, $\omega$
$\theta_{\max}$ , completeness:	25.0°, 94%
$N(hkl)_{\text{measured}}$ , $N(hkl)_{\text{unique}}$ , $R_{\text{int}}$ :	5684, 2860, 0.030
Criterion for $I_{\text{obs}}$ , $N(hkl)_{\text{gt}}$ :	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$ , 2487
$N(\text{param})_{\text{refined}}$ :	148
Programs:	Diamond [1], IPDS [2], SHELX [3, 4]

## Source of material

Iodine ( $\geq 99.0\%$ ), cesium iodide, 1,4,7,10-tetraoxacyclododecan (12-crown-4) and ethanol were purchased from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). All reagents were of analytical grade and used as purchased. The title compound was prepared by dissolving 0.16 g (0.63 mmol) of CsI and 0.48 g (1.89 mmol)  $I_2$  in a 20 mL ethanol/10 mL methanol mixture at room temperature. Then 0.2 mL (1.26 mmol) 12-crown-4 is added while stirring. The dark, clear solution precipitates after 5 days at room temperature black crystals of  $[Cs(12\text{-crown-4})_2]I_5$ .

## Experimental details

A black single crystal was chosen from the mother liquor and measured at 293 K on an IPDS diffractometer [2]. The structure solution, refinement and other calculations were done with the programs SHELXL [3, 4]. All antimicrobial studies were performed as described in our previous work [5].

## Comment

Iodine has important applications as antimicrobial agent because of its well known biocidal properties [6, 7]. Iodine and related compounds in form of nanoparticles are already used in wound dressing applications [8]. The drawbacks of iodine use in wound care are sublimation and skin irritation [6, 8]. The iodine content can be stabilized by designing polyiodides complexed by crown ethers as previously reported in

<https://doi.org/10.1515/ncrs-2020-0057>

Received January 29, 2020; accepted February 18, 2020; available online February 28, 2020

## Abstract

$C_{16}H_{32}CsI_5O_8$ , monoclinic,  $I2/a$  (no. 15),  $a = 16.386(2)$  Å,  $b = 11.7050(10)$  Å,  $c = 19.005(3)$  Å,  $\beta = 108.008(14)^\circ$ ,  $V = 3466.6(8)$  Å<sup>3</sup>,  $Z = 4$ ,  $R_{\text{gt}}(F) = 0.0437$ ,  $wR_{\text{ref}}(F^2) = 0.1085$ ,  $T = 293(2)$  K.

CCDC no.: 1984799

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of

\*Corresponding author: Zehra Edis, College of Pharmacy and Health Sciences, Ajman University, PO Box 346, Ajman, UAE, e-mail: z.edis@ajman.ac.ae. <https://orcid.org/0000-0001-9555-3473>

Samir Haj Bloukh: College of Pharmacy and Health Sciences, Ajman University, PO Box 346, Ajman, UAE

**Table 2:** Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>).

Atom	x	y	z	$U_{iso}^*/U_{eq}$
Cs	0.250000	0.32510(5)	0.500000	0.0404(2)
I1	0.250000	0.70462(6)	0.500000	0.0437(2)
I2	0.24493(3)	0.70796(5)	0.34433(3)	0.0509(2)
I3	0.07725(4)	0.51639(5)	0.27633(3)	0.0516(2)
I4	-0.06144(4)	0.37426(4)	0.22230(2)	0.0508(2)
O1	0.4132(4)	0.1738(6)	0.5618(3)	0.0622(16)
O2	0.4332(4)	0.3771(5)	0.4876(3)	0.0554(13)
O3	0.3138(4)	0.3016(5)	0.3545(3)	0.0559(15)
O4	0.2921(4)	0.0959(5)	0.4279(4)	0.0620(15)
C1	0.4931(7)	0.2045(10)	0.5599(5)	0.076(3)
H1A	0.504180	0.171193	0.517043	0.092*
H1B	0.536615	0.177647	0.604045	0.092*
C2	0.4951(6)	0.3297(9)	0.5560(5)	0.066(3)
H2A	0.552593	0.353692	0.558440	0.079*
H2B	0.482476	0.361265	0.598651	0.079*
C3	0.4593(6)	0.3638(9)	0.4255(4)	0.062(2)
H3A	0.480532	0.286993	0.423253	0.074*
H3B	0.504519	0.417736	0.426237	0.074*
C4	0.3826(6)	0.3858(7)	0.3617(4)	0.054(2)
H4A	0.360721	0.461366	0.366748	0.065*
H4B	0.398926	0.385145	0.316819	0.065*
C5	0.3286(6)	0.1971(7)	0.3276(4)	0.054(2)
H5A	0.388029	0.174850	0.349861	0.065*
H5B	0.316729	0.200859	0.274391	0.065*
C6	0.2711(6)	0.1133(8)	0.3465(4)	0.055(2)
H6A	0.275470	0.040790	0.323199	0.066*
H6B	0.212256	0.139616	0.326735	0.066*
C7	0.3660(6)	0.0329(8)	0.4599(5)	0.067(2)
H7A	0.411653	0.058069	0.440996	0.080*
H7B	0.355311	-0.047501	0.448353	0.080*
C8	0.3903(6)	0.0506(7)	0.5391(5)	0.065(3)
H8A	0.439129	0.002333	0.562963	0.078*
H8B	0.343159	0.027207	0.556632	0.078*

our group [9, 10]. Triiodides with halogen bonding are usually stable compounds with many potential applications [11–13]. Triiodides with halogen bonding can be used as antimicrobial agents because they are stable and gradually release free iodine [5]. In these compounds, the released free iodine interacts with cell membranes and leads to protein oxidation [5, 6, 8].

Pentaiodides may have also microbicidal properties and are generally formed by combination of iodine and triiodide units [14–16]. Many interesting compounds have been reported so far [9, 17, 18]. We synthesized the cesium-12-crown-4 polyiodide and redetermined its structure as a penta-iodide. The two crown ether molecules form a sandwich complex around the cation like in our previous compounds [5, 9, 10]. The anionic structure consists of a remarkable I<sub>5</sub><sup>-</sup> polymer, which has similarity to the anionic structure in nicotine-1,1'-diumbis(triiodide)-diiodine (1/1) [16].

In our title compound [Cs(12-crown-4)<sub>2</sub>]I<sub>5</sub>, the asymmetric unit contains symmetrical, linear triiodide anions I<sub>3</sub><sup>-</sup> with crystallographic symmetry of a 2-fold axis. Two iodine

molecules (I3-I4) are weakly connected to these symmetric triiodide (I2-I1-I2') halogen bond donor. The resulting penta-iodide structure is a zig-zag chain. The bond lengths and angles within this structure is in expected ranges [16], which is also true for the cationic Cs(I) complex [9] and are further examples of a three-center-system [I-I-I]<sup>-</sup> with halogen bonding as our previously reported triiodides [5, 9, 10]. This compound is the ideal candidate to act as an antimicrobial agent. Therefore we investigated its inhibitory effects on common microorganisms.

[Cs(12-crown-4)<sub>2</sub>]I<sub>5</sub> exhibited antimicrobial activity against several pathogens. The disc diffusion and agar well assays revealed strong antimicrobial activity against the reference strains *Candida albicans* WDCM 00054 (zone of inhibition ZOI = 47 mm), *Escherichia coli* WDCM 00013 (ZOI = 15 mm), *Streptococcus pyogenes* ATCC 19615 (ZOI = 25 mm) and the clinical sample *Streptococcus pneumoniae* (ZOI = 26 mm). Our previous compound [Na(12-crown-4)<sub>2</sub>]I<sub>3</sub> acted as a strong antifungal and broad-spectrum bacteriocidal agent [5]. In comparison, the same reference strain *C. albicans* WDCM 00054 was inhibited less strongly (ZOI = 40 mm). The Gram negative *E. coli* was in comparison to [Cs(12-crown-4)<sub>2</sub>]I<sub>5</sub> less resistant (ZOI = 23 mm) to the triiodide [5]. The Gram positive bacteria *S. pyogenes* showed a larger inhibition zone of 34 mm in [Na(12-crown-4)<sub>2</sub>]I<sub>3</sub>, while *S. pneumoniae* (ZOI = 28 mm) exhibited a slightly stronger result than in the penta-iodide [Cs(12-crown-4)<sub>2</sub>]I<sub>5</sub>.

These results support our hypothesis, that polyiodides can be used as antimicrobial agents. Once the complex compound interacts with the microbial cell membranes by electrostatic interactions, the title compound decomposes and results in controlled free molecular iodine release [5].

**Acknowledgements:** We are grateful to Ajman University, Ajman, UAE for funding this research by the Ajman University-IRG Research Grant (2019-IRG-PH-5). We express our deep gratitude to Guido J. Reiss for kindly managing our structural data.

## References

1. Brandenburg, K.: DIAMOND. Visual Crystal Structure Information System. Ver. 4.0. Crystal Impact, Bonn, Germany (2015).
2. Stoe & Cie. IPDS Software. Version 2.90. Stoe & Cie, Darmstadt, Germany (1999).
3. Sheldrick, G. M.: SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr.* **A71** (2015) 3–8.
4. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
5. Edis, Z.; Haj Bloukh, S.; Abu Sara, H.; Bhakhoa, H.; Rhyman, L.; Ramasami, P.: “Smart” triiodide compounds: does halogen

- bonding influence antimicrobial activities?. *Pathogens* **8** (2019) 182–202.
- Kaiho, T.: Iodine chemistry and applications. (Ed.: T. Kaiho), John Wiley & Sons, Inc.: Hoboken, NJ, USA (2015), ISBN: 978-1-118-46629-2.
  - He, C.; Parrish, D. A.; Shreeve, J. M.: Alkyl ammonium cation stabilized biocidal polyiodides with adaptable high density and low pressure. *Chem. Eur. J.* **20** (2014) 6699–6706.
  - Viswanathan, K.; Babu, D. B.; Jayakumar, G.; Raj, G. D.: Antimicrobial and skin wound dressing application of molecular iodine nanoparticles. *Mater. Res. Express* **4** (2017) 104003.
  - Edis, Z.; Haj Bloukh, S.: Preparation, structural and spectroscopic characterization of a pentaiodide  $[Rb(12\text{-Crown-}4)_2]I_5$ . *Z. Naturforsch.* **68b** (2013) 1340–1346.
  - Edis, Z.; Haj Bloukh, S.: Preparation and structural and spectroscopic characterization of triiodides  $[M(12\text{-Crown-}4)_2]I_3$  with  $M = Na$  and  $Rb$ . *Z. Naturforsch.* **69b** (2014) 995–1002.
  - Bartashevich, E. V.; Grigoreva, E. A.; Yushina, I. D.; Bulatova, L. M.; Tsirelson, V. G.: Modern level for the prediction of properties of iodine-containing organic compounds: iodine forming halogen bonds. *Russ. Chem. Bull. Int. Ed.* **66** (2017) 1–12.
  - Reiss, G. J.: Halogen and hydrogen bonding in the layered crystal structure of 2-iodoanilinium triiodide,  $C_6H_7I_4N$ . *Z. Kristallogr. NCS* **234** (2019) 899–902.
  - van Megen, M.; Reiss, G. J.:  $I_6^{2-}$  Anion composed of two asymmetric triiodide moieties: a competition between halogen and hydrogen bond. *Inorganics* **1** (2013) 3–13 and cited articles.
  - Blake, A. J.; Devillanova, F. A.; Gould, R. O.; Li, W. S.; Lippolis, V.; Parsons, S.; Radek, C.; Schröder, M.: Template selfassembly of polyiodide networks. *Chem. Soc. Rev.* **27** (1998) 195–205.
  - Svensson, P. H.; Kloo, L.: Synthesis, structure, and bonding in polyiodide and metal iodide-iodine systems. *Chem. Rev.* **103** (2003) 1649–1684.
  - Reiss, G. J.:  $I_5^-$  polymers with a layered arrangement: synthesis, spectroscopy, and structure of a new polyiodide salt in the nicotine/HI/I<sub>2</sub> system. *Z. Naturforsch.* **B70** (2015) 735–739.
  - Walbaum, C.; Pantenburg, I.; Meyer, G.: Penta-, Hepta- und Oктаiodid-Anionen in Salzen mit Erdalkalimetall-Kronenether-Kationen. *Z. Naturforsch.* **B65** (2010) 1077–1083.
  - Reiss, G. J.: A cyclic  $I_{10}^{2-}$  anion in the layered crystal structure of theophyllinium pentaiodide,  $C_7H_9I_5N_4O_2$ . *Z. Kristallogr. NCS* **234** (2019) 737–739.