



Alexandria University  
**Alexandria Engineering Journal**

[www.elsevier.com/locate/aej](http://www.elsevier.com/locate/aej)  
[www.sciencedirect.com](http://www.sciencedirect.com)



# Numerical schemes for studying biomathematics model inherited with memory-time and delay-time



Imad Jaradat<sup>a,\*</sup>, Marwan Alquran<sup>a</sup>, Shaher Momani<sup>b,c</sup>, Dumitru Baleanu<sup>d,e</sup>

<sup>a</sup> Department of Mathematics and Statistics, Jordan University of Science and Technology, P.O. Box: 3030, Irbid 22110, Jordan

<sup>b</sup> Department of Mathematics and Sciences, College of Humanities and Sciences, Ajman University, Ajman, United Arab Emirates

<sup>c</sup> Department of Mathematics, Faculty of Science, The University of Jordan, Amman 11942, Jordan

<sup>d</sup> Department of Mathematics, Cankaya University, Ankara, Turkey

<sup>e</sup> Institute of Space Sciences, Magurele, Bucharest, Romania

Received 29 January 2020; revised 3 March 2020; accepted 11 March 2020

Available online 15 May 2020

## KEYWORDS

Caputo-time-fractional derivative;  
 Delay time;  
 Fractional power series;  
 Homotopy perturbation method;  
 Mathematical biological model

## MSC(2010)

35F25;  
 35C10

**Abstract** The effect of inherited memory-time and delay-time in the formulation of a mathematical population growth model is considered. Two different numerical schemes are introduced to study analytically the propagation of population growth. We provide a graphical analysis that shows the impact of both memory-time and delay-time acting on the behavior of population density. We concluded that both delay-time and time-fractional-derivative play the same role as delaying the propagation of the nonlinear population growth.

© 2020 The Authors. Published by Elsevier B.V. on behalf of Faculty of Engineering, Alexandria University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

A variety of nonlinear evolution equations that represent real-life applications are exposure for unexpected physical changes such as, delaying in its propagations, bifurcating its solution or stability issues. Linking such change with the theoretical considerations may lead to unknown. A wise thought on reasons

behind such unexpected bifurcations is directed to the role of the rate of change of the field function with respect to time. Many studies emerged to review the rate of change as a fractional hospital with a topological relationship with the integer case. All scientific researches become interested in this area taking into consideration the use of the fractional derivative instead of the integer-derivative [1–9]. Extensive works have been accomplished to draw suitable mathematical methods for dealing with these fractal issues. Among the most important and widely used numerical methods are: decomposition schemes [10,11], variational iteration schema [12,13], fractional power series methods [14–17], finite difference scheme [18], Sumudu decomposition method [19] and other methods [20–22].

\* Corresponding author.

E-mail addresses: [iajaradat@just.edu.jo](mailto:iajaradat@just.edu.jo) (I. Jaradat), [marwan04@just.edu.jo](mailto:marwan04@just.edu.jo) (M. Alquran), [s.momani@ju.edu.jo](mailto:s.momani@ju.edu.jo) (S. Momani), [dumitru@cankaya.edu.tr](mailto:dumitru@cankaya.edu.tr) (D. Baleanu).

Peer review under responsibility of Faculty of Engineering, Alexandria University.

<https://doi.org/10.1016/j.aej.2020.03.038>

1110-0168 © 2020 The Authors. Published by Elsevier B.V. on behalf of Faculty of Engineering, Alexandria University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

One of the interesting mathematical models arise in science is the biology population growth model which is governed by the following form [23,24]

$$D_t P(x, t) - D_{xx}(P^2(x, t)) - f(P(x, t)) = 0, \quad t > 0. \tag{1.1}$$

The function  $P(x, t)$  stands for the population density and the functional  $f(P)$  is the reaction source that describes the dynamics of the population growth. There are different types of the reaction source;  $f(P) = cP$ , is identified by Malthusian law.  $f(P) = aP - bP^2$ , is regarded as the Verhulst law. The most general form of such reaction source is defined as  $f(P) = aP^n(1 - \frac{b}{a}P^n)$ .

The main goal of this work is to study the above model under new considerations; the time-integer-derivative is to be replaced by fractional order and the reaction source is of Verhulst-type acting with delay type of form  $\tau t$ :  $0 < \tau < 1$ . Therefore, the new biological model has the form:

$$D_t^\alpha P(x, t) - D_{xx}(P^2(x, t)) - aP(x, \tau t) + bP^2(x, \tau t) = 0, \quad t > 0, \tag{1.2}$$

where  $0 < \alpha \leq 1$  is the Caputo-fractional derivative [25–27].

The model given in (1.2) is to be presented for the first time. Whereas, previous studies concerned with only time-fractional derivatives. For example, in [28], the conformable fractional-derivative is used and exact solutions are found by means of Kudryashov method. Two-dimensional form of the time-fractional population growth of Malthusian law [29] is studied by means of fractional variational iteration method (FVIM). Finally, The finite difference method (FDM) and variational iteration method (VIM) have been successfully implemented [30] for solving (1.2) but with the absence of delay action.

The format of the current paper is as follows. In Section 2, we provide in details the t-spectrum scheme to solve the proposed model and give some graphical analysis. In Section 3, we present the homotopy perturbation method as an alternative scheme to handle (1.2). Finally, some concluding remarks and observations are given in the last section.

## 2. Time-coordinate spectrum function method

A power series of the form [31,32]

$$\sum_{j=0}^{\infty} b_j t^{j\alpha} = b_0 + b_1 t^\alpha + b_2 t^{2\alpha} + \dots, \tag{2.1}$$

is called fractional Maclaurin series (FMS). Suppose that the function  $v(t)$  has FMS of the form

$$v(t) = \sum_{j=0}^{\infty} b_j t^{j\alpha}, \quad 0 < t < R, \tag{2.2}$$

if  $D^{j\alpha} v(t)$  are continuous on  $I = (0, R)$ , then  $b_j = \frac{D^{j\alpha} v(0)}{\Gamma(j\alpha+1)}$  and  $R$  is called the radius of convergence.

Consider that the solution of (1.2) can be represented as a product of two single variable functions,  $P(x, t) = u(x)v(t)$ , where  $u(x)$  is analytic function and  $v(t)$  has FMS representation. Then,

$$P(x, t) = \left( \sum_{i=0}^{\infty} a_i x^i \right) \left( \sum_{j=0}^{\infty} b_j t^{j\alpha} \right), = \sum_{k=0}^{\infty} U_k(x) t^{k\alpha}. \tag{2.3}$$

where  $U_k(x) = \frac{D^{k\alpha} v(x,0)}{\Gamma(k\alpha+1)}$ . We refer to  $U_k(x)$  as the transformed function or the t-coordinate spectrum function. Also, the inverse of the spectrum  $U_k(x)$  is given by (2.3). Apply the t-fractional derivative on (2.3) gives

$$D_t^\alpha P(x, t) = \sum_{k=0}^{\infty} \left( \frac{\Gamma((k+1)\alpha+1)}{\Gamma(k\alpha+1)} U_{k+1}(x) \right) t^{k\alpha}. \tag{2.4}$$

Also,

$$P(x, \tau t) = \sum_{k=0}^{\infty} (t^{k\alpha} U_k(x)) t^{k\alpha}. \tag{2.5}$$

Apply the cauchy product of two infinite series and by (2.5) we get the following expansion

$$P^2(x, \tau t) = \sum_{k=0}^{\infty} \left( t^{k\alpha} \sum_{n=0}^k U_{k-n}(x) U_n(x) \right) t^{k\alpha}. \tag{2.6}$$

Table 1, provides the necessary transformed spectrum functions that will be used to solve the problem under investigation. Applying the rules given in Table 1, the following spectrum transformed form of Eq. (1.2) is obtained

$$\frac{\Gamma((k+1)\alpha+1)}{\Gamma(k\alpha+1)} U_{k+1}(x) - \sum_{n=0}^k (U_n(x) U_{k-n}(x))'' - a\tau^{k\alpha} U_k(x) + b\tau^{k\alpha} \sum_{n=0}^k U_n(x) U_{k-n}(x) = 0. \tag{2.7}$$

From (2.7), we deduce the recurrence relation regarding  $U_k(x)$  as

$$U_{k+1}(x) = \frac{\Gamma(k\alpha+1)}{\Gamma((k+1)\alpha+1)} \left( \sum_{n=0}^k (U_n(x) U_{k-n}(x))'' + a\tau^{k\alpha} U_k(x) - b\tau^{k\alpha} \sum_{n=0}^k U_n(x) U_{k-n}(x) \right). \tag{2.8}$$

For clarity, we list the first few terms of the spectrum sequence of the growth model

$$\begin{aligned} U_1 &= \frac{1}{\Gamma(\alpha+1)} \left( (U_0'') + aU_0 - bU_0^2 \right), \\ U_2 &= \frac{\Gamma(\alpha+1)}{\Gamma(2\alpha+1)} \left( 2(U_0 U_1)'' + a\tau^\alpha U_1 - 2b\tau^\alpha U_0 U_1 \right), \\ U_3 &= \frac{\Gamma(2\alpha+1)}{\Gamma(3\alpha+1)} \left( 2(U_0 U_2)'' + (U_1'') + a\tau^{2\alpha} U_2 - b\tau^{2\alpha} (2U_0 U_2 + U_1^2) \right), \\ U_4 &= \frac{\Gamma(3\alpha+1)}{\Gamma(4\alpha+1)} \left( 2(U_0 U_3)'' + 2(U_1 U_2)'' + a\tau^{3\alpha} U_3 - 2b\tau^{3\alpha} (U_0 U_3 + U_1 U_2) \right). \end{aligned} \tag{2.9}$$

Now, we are ready to discuss some graphical analysis regarding the accuracy of t-spectrum function method, and the impact of both time-fractional and time-delay acting on the

**Table 1** Transformed t-coordinate spectrum functions.

$P(x, t)$	$U_k(x)$
$P(x, \tau t)$	$t^{k\alpha} U_k(x)$
$P^2(x, t)$	$\sum_{n=0}^k U_{(k-n)}(x) U_n(x)$
$P^2(x, \tau t)$	$\tau^{k\alpha} \sum_{n=0}^k U_{(k-n)}(x) U_n(x)$
$D_t^\alpha P(x, t)$	$\frac{\Gamma((k+1)\alpha+1)}{\Gamma(k\alpha+1)} U_{k+1}(x)$
$D_{xx} P^2(x, t)$	$\sum_{n=0}^k (U_{(k-n)}(x) U_n(x))''$

population density. The analysis will be performed on a numerical example of (1.2); the case of  $a = b = 1$ ; subject to the initial population density  $P(x, 0) = f(x) = 1 + e^{\frac{x}{2}}$ . We should point here that the presence of the delay factor  $\tau$  in the calculations of the spectrum functions  $U_i(x)$  will increase the size of the resulting terms. Moreover, it is hard to find or even guess a closed form solution. Therefore, we decide to consider  $\phi = \phi(x, t) = \sum_{n=0}^6 U_n(x)t^{n\alpha}$ , to represent the approximation solution for the population problem. Fig. 1, presents, respectively, the exact solution  $P(x, t) = 1 + e^{\frac{x-t}{2}}$  (See [28]), the approximate function  $\phi$ , and the obtained absolute error  $|P(x, t) - \phi(x, t)|$  under the case of  $\alpha = \tau = 1$ . These obtained 3D plots show that the assigned approximation expansion is in good agreement with the exact one. Fig. 2, shows the impact and the interaction of both fractional order  $\alpha$  and the delay factor  $\tau$  on the population density. Four drawn relations can be observed:

- The order of  $\alpha$  :  $0 < \alpha < 1$  is proportional with the population density.
- An increase of the delay factor leads to an increase in the population density. Moreover, for any value of  $\tau$  :  $0 < \tau < 1$ , it delays the expected value of the population growth.
- In the presence of the fractional-derivative “ $\alpha < 1$ ”, the delay factor affects the delay of population growth much faster if compared with the integer-case “ $\alpha = 1$ ”.
- Both delay-time and time-fractional-derivative play the same role as delaying the propagation of nonlinear evolutionary models.

In conclusion, we speculate that the fractional derivative order or/and the time-delay have the potential to affect real-life observation of the dynamics of interacting populations.

2.1. Convergence of spectrum function method

In this part, we validate numerically the convergence of the transformed t-spectrum function method applied to the previous example. Therefore, to study the convergence of the obtained approximate solution  $\phi(x, t)$  to its exact solution, we use the approach proposed by Hosseini and Nasabzadeh [33]. We determine the values of  $\gamma_i$  where  $\gamma_i = \max\{\frac{U_{i+1}(x)}{U_i(x)}\} : x \in (-\infty, \infty)$  and therefore, the solution converges when  $|\gamma_i| < 1$ . For the proposed example, we obtained the following bounds:

- $-\frac{1}{2} < \gamma_0 < 0$ .
- $\gamma_1 = -\frac{1}{4}$ .
- $\gamma_2 = -\frac{1}{6}$ .
- $\gamma_3 = -\frac{1}{8}$ .
- $\gamma_4 = -\frac{1}{10}$ .
- $\gamma_5 = -\frac{1}{12}$ .
- $\gamma_k = -\frac{1}{2(k+1)} : k \geq 1$ .

The above findings are regarded as a numerical evidence that the solution  $\phi(x, t)$  obtained by using transformed spectrum method converges to  $P(x, t)$ .





Fig. 1 3D plots of, respectively,  $P(x, t)$ ,  $\phi(x, t)$  and  $|P(x, t) - \phi(x, t)|$ . The case for  $\alpha = \tau = 1$  and  $n = 6$ .



**Fig. 2** The interaction of both fractional order  $\alpha$  and the delay factor  $\tau$  on the population density. Where, Red  $\equiv (\tau = 1)$ , Black  $\equiv (\tau = 0.75)$ , Blue  $\equiv (\tau = 0.5)$ , Green  $\equiv (\tau = 0.25)$ .

**3. Homotopy perturbation schema**

The Homotopy perturbation method (HPM) will be used as an alternative technique to solve (1.2). This numerical scheme suggest to write the problem in two homotopy forms [34,35]; either

$$D_t P(x, t) = q(D_t P(x, t) + D_t^\alpha P(x, t) - D_{xx}(P^2(x, t)) - aP(x, \tau t) + bP^2(x, \tau t)), \tag{3.1}$$

or

$$D_t^\alpha P(x, t) = q(D_{xx}(P^2(x, t)) + aP(x, \tau t) - bP^2(x, \tau t)), \tag{3.2}$$

with a perturbation form of the function  $P(x, t)$  as

$$P(x, t) = \sum_{i=0}^{\infty} q^i P_i(x, t). \tag{3.3}$$

To solve (1.2) subject to  $P(x, 0) = f(x)$ , we substitute (3.3) in either (3.1) or (3.2) and in the initial condition. Then, in a sequence order, we solve sub-equations resulting from collecting same coefficients of  $q^i$ . By considering (3.1) we use the rule  $D_t^\alpha t^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta-\alpha+1)} t^{\beta-\alpha}$  whereas in the case of (3.2) we use the rule  $J_t^\alpha t^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta+\alpha+1)} t^{\beta+\alpha}$ . In this context, we prefer the form given in (3.2). Now, we proceed as the following steps:

**Step I:**  $q^0$ : We solve

$$D_t^0 P_0(x, t) = 0, \quad P_0(x, 0) = f(x).$$

Which leads to the first component

$$P_0(x, t) = f(x). \tag{3.4}$$

**Step II:**  $q^1$ : We solve

$$D_t^1 P_1(x, t) = D_{xx}(P_0^2(x, t)) + aP_0(x, \tau t) - bP_0^2(x, \tau t), \quad P_1(x, 0) = 0.$$

Which leads to the second component

$$P_1(x, t) = h(x) \frac{t^\alpha}{\Gamma(\alpha + 1)}, \tag{3.5}$$

where  $h(x) = f''(x) + af(x) - bf^2(x)$ .

**Step III:**  $q^2$ : We solve

$$D_t^2 P_2(x, t) = D_{xx}(2P_0(x, t)P_1(x, t)) + aP_1(x, \tau t) - 2bP_0(x, \tau t)P_1(x, \tau t), \quad P_2(x, 0) = 0.$$

Which leads to the third component

$$P_2(x, t) = \{2(f(x)h(x))'' + (a - 2bf(x))h(x)\tau^\alpha\} \times \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}. \tag{3.6}$$


**Step IV:**  $q^3$ : We solve

$$D_t^3 P_3(x, t) = D_{xx}(2P_0(x, t)P_2(x, t) + P_1^2(x, t)) + aP_2(x, \tau t) - b(2P_0(x, \tau t)P_2(x, \tau t) + P_1^2(x, \tau t)), \quad P_3(x, 0) = 0.$$

Which leads to the fourth component

$$P_3(x, t) = \{2f(x)(2(f(x)h(x))'' + (a - 2bf(x))h(x)\tau^\alpha)\}'' + \frac{\Gamma(2\alpha + 1)}{\Gamma^2(\alpha + 1)} (h^2(x))'' \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} + \{a(2(f(x)h(x))'' + (a - 2bf(x))h(x)\tau^\alpha)\tau^{2\alpha}\} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} - b\{2f(x)(2(f(x)h(x))'' + (a - 2bf(x))h(x)\tau^\alpha)\} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} + h^2(x) \frac{\Gamma(2\alpha + 1)}{\Gamma^2(\alpha + 1)} \tau^{2\alpha} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)}. \tag{3.7}$$

We observe the difficulty in determining the components  $P_i(x, t)$  as we proceed further for  $i \geq 3$ . To this end, we con-



**Fig. 3** 3D plots of, respectively,  $P(x, t)$ ,  $\psi(x, t)$  and  $|P(x, t) - \psi(x, t)|$ . The case for  $\alpha = \tau = 1$  and  $n = 3$ .

sider  $\psi(x, t) = \sum_{i=0}^3 P_i(x, t)$  to represent the homotopy approximate solution to (1.2). Fig. 3, shows that HPM is a good alternative scheme even considering a few components of its series' representation.

**4. Conclusions**

In this manuscript, we suggested a new version of the mathematical biology growth model. The new model consider the presence of two actions on the time-coordinate. These actions are categorized as fractional-time derivative to represent the rate of change in the population density, and a delay-time of pantograph type acting on the reaction source term which is responsible for the dynamics of the population growth.

We offered two numerical schemes to find an approximation solution of the population growth model. The first method is derived from the fractional powers series whose coefficients are addressed as the time-coordinate spectrum functions. This approach leads to a recursive relation among its spectrum functions which give the advantages of determining "as many as needed" the spectrum functions with delicate computational process and in turns leads to a high accuracy approximation. In contrast, the alternative homotopy perturbation technique which decomposes the solution as infinite analytical series, its components to be determined by solving successive initial value fractional problems. These homotopy steps become difficult task as the calculation processes go farther. In most cases, only a few terms are considered for the approximate solution with limited accuracy size.

The obtained findings for the t-spectrum function method are used to study the impact of both time-fractional derivative

and the delay-time. We concluded that the presence of fractional derivative helps to understand the fact that the delay occurs in the population density "due to a delay-factor embedded in the problem" could be less than the expected values. We may say that integer-derivative can be regarded as theoretical-perspective, whereas the fractional-derivative is a realistic-perspective.

As future work, we will considered the n-dimensional extension of the fractional power series [36–40] to study multivariate-fractional models arise in sciences and engineering.

**Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Acknowledgements**

The first and the third authors would like to thank Ajman University, Ajman, UAE for providing the financial support.

**References**

- [1] A. Atangana, D. Baleanu, Caputo-Fabrizio derivative applied to groundwater flow within confined aquifer, *J. Eng. Mech.* 143 (5) (2017) D4016005.
- [2] K.M. Saad, D. Baleanu, A. Atangana, New fractional derivatives applied to the Korteweg-de Vries and Korteweg-de Vries-Burger's equations, *Comput. Appl. Math.* 37 (4) (2018) 5203–5216.

- [3] K.M. Saad, A. Atangana, D. Baleanu, New fractional derivatives with non-singular kernel applied to the Burgers equation, *CHAOS* 28 (6) (2018) 063109.
- [4] A. Atangana, D. Baleanu, New fractional derivatives with non-local and non-singular kernel Theory and Application to Heat Transfer Model, *Therm. Sci.* 20 (2) (2016) 763–769.
- [5] A. Jaradat, M.S.M. Noorani, M. Alquran, H.M. Jaradat, A Novel method for solving Caputo-time-fractional dispersive long Wave Wu-Zhang system, *Nonlinear Dyn. Syst.Theory* 18 (2) (2018) 182–190.
- [6] M. Alquran, I. Jaradat, A novel scheme for solving Caputo time-fractional nonlinear equations: theory and application, *Nonlinear Dyn.* 91 (4) (2018) 2389–2395.
- [7] J. Singh, D. Kumar, D. Baleanu, On the analysis of fractional diabetes model with exponential law, *Adv. Differ. Eqs.* 2018 (2018) 231.
- [8] D. Kumar, J. Singh, K. Tanwar, D. Baleanu, A new fractional exothermic reactions model having constant heat source in porous media with power, exponential and Mittag-Leffler Laws, *Int. J. Heat Mass Transf.* 138 (2019) 1222–1227.
- [9] J. Singh, A. Kilicman, D. Kumar, R. Swroop, Md. Ali Fadzilah, Numerical study for fractional model of nonlinear predator-prey biological population dynamical system, *Therm. Sci.* 23 (6) (2019) 2017–2025.
- [10] K. Al-Khaled, F. Allan, Decomposition method for solving nonlinear integro-differential equations, *J. Appl. Math. Comput.* 19 (1–2) (2005) 415.
- [11] E.A. Az-Zo'bi, K. Al-Khaled, A new convergence proof of the Adomian decomposition method for a mixed hyperbolic elliptic system of conservation laws, *Appl. Math. Comput.* 217 (8) (2010) 4248–4256.
- [12] A.M. Wazwaz, The variational iteration method: a reliable analytic tool for solving linear and nonlinear wave equations, *Comput. Math. Appl.* 54 (7–8) (2007) 926–932.
- [13] J.H. He, Variational iteration method-Some recent results and new interpretations, *J. Comput. Appl. Math.* 207 (1) (2007) 3–17.
- [14] H.M. Jaradat, I. Jaradat, M. Alquran, M.M.M. Jaradat, Z. Mustafa, K. Abohassan, R. Abdelkarim, Approximate solutions to the generalized time-fractional Ito system, *Italian J. Pure Appl. Math.* 37 (2017) 699–710.
- [15] M. Alquran, I. Jaradat, S. Sivasundaram, Elegant scheme for solving Caputo-time-fractional integro-differential equations, *Nonlinear Stud.* 25 (2) (2018) 385–393.
- [16] I. Jaradat, M. Al-Dolat, K. Al-Zoubi, M. Alquran, Theory and applications of a more general form for fractional power series expansion, *Chaos Solitons Fract.* 108 (2018) 107–110.
- [17] I. Jaradat, M. Alquran, M. Al-Dolat, Analytic solution of homogeneous time-invariant fractional IVP, *Adv. Differ. Eqs.* 2018 (2018) 143.
- [18] C. Li, F. Zeng, The finite difference methods for fractional ordinary differential equations, *Numer. Funct. Anal. Optim.* 34 (2) (2013) 149–179.
- [19] K. Al-Khaled, Numerical solution of time-fractional partial differential equations using Sumudu decomposition method, *Rom. J. Phys* 60 (1–2) (2015) 99–110.
- [20] D. Kumar, J. Singh, S.D. Purohit, R. Swroop, A hybrid analytical algorithm for nonlinear fractional wave-like equations, *Math. Modell. Nat. Phenomena* 14 (2019) 304.
- [21] S. Bhattar, A. Mathur, D. Kumar, K.S. Nisar, J. Singh, Fractional modified Kawahara equation with Mittag-Leffler law, *Chaos Solitons Fract.* 131 (2020) 109508.
- [22] A. Goswami, J. Singh, D. Kumar, Sushila, An efficient analytical approach for fractional equal width equations describing hydro-magnetic waves in cold plasma, *Physica A* 524 (2019) 563–575.
- [23] K.T. Alligood, T.D. Sauer, J.A. Yorke, *An Introduction to Dynamical Systems*, Springer, 1996.
- [24] J.M. Cushing, *An Introduction to Structured Population Dynamics*, Society for Industrial and Applied Mathematics, 1998.
- [25] G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, Yverdon, 1993.
- [26] F. Mainardi, G. Panini, The Wright functions as solutions of the time-fractional diffusion equation, *Appl. Math. Comput.* 141 (2003) 51–62.
- [27] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [28] Kumar D. Dipankar, A.R. Seadawy, A.K. Joardare, Modified Kudryashov method via new exact solutions for some conformable fractional differential equations arising in mathematical biology, *Chinese J. Phys.* 56 (1) (2018) 75–85.
- [29] A. Prakash, M. Kumar, Numerical solution of two dimensional time fractional-order biological population model, *Open Phys.* 14 (2016) 177–186.
- [30] N.H. Sweilam, M.M. Khader, A.M.S. Mahdy, Numerical studies for solving fractional-order logistic equation, *Int. J. Pure Appl. Math.* 78 (8) (2012) 1199–1210.
- [31] Arqub O. Abu, Series solution of fuzzy differential equations under strongly generalized differentiability, *J. Adv. Res. Appl. Math.* 5 (2013) 31–52.
- [32] Arqub O. Abu, A. El-Ajou, Zhou Z. Al, S. Momani, Multiple solutions of nonlinear boundary value problems of fractional order: a new analytic iterative technique, *Entropy* 16 (2014) 471–493.
- [33] M.M. Hosseini, H. Nasabzadeh, On the convergence of Adomian decomposition method, *Appl. Math. Comput.* 182 (2006) 536–543.
- [34] M. Ali, M. Alquran, M. Mohammad, Solitonic solutions for homogeneous KdV systems by homotopy analysis method, *J. Appl. Math.* 2012 (2012) 569098.
- [35] M. Alquran, M. Mohammad, Approximate solutions to system of nonlinear partial differential equations using homotopy perturbation method, *Int. J. Nonlinear Sci.* 12 (4) (2011) 485–497.
- [36] I. Jaradat, M. Alquran, R. Abdel-Muhsen, An analytical framework of 2D diffusion, wave-like, telegraph, and Burgers' models with twofold Caputo derivatives ordering, *Nonlinear Dyn.* 93 (4) (2018) 1911–1922.
- [37] M. Alquran, I. Jaradat, D. Baleanu, R. Abdel-Muhsen, An analytical study of  $(2+1)$ -dimensional physical models embedded entirely in fractal space, *Rom. J. Phys.* 64 (2019) 103.
- [38] M. Alquran, I. Jaradat, R. Abdel-Muhsen, Embedding  $(3+1)$ -dimensional diffusion, telegraph, and Burgers' equations into fractal 2D and 3D spaces: an analytical study, *J. King Saud Univ.-Sci.* 32 (1) (2020) 349–355.
- [39] I. Jaradat, M. Alquran, F. Yousef, S. Momani, D. Baleanu, On  $(2+1)$ -dimensional physical models endowed with decoupled spatial and temporal memory indices, *Eur. Phys. J. Plus* 134 (7) (2019) 360.
- [40] F. Yousef, M. Alquran, I. Jaradat, S. Momani, D. Baleanu, New fractional analytical study of three-dimensional evolution equation equipped with three memory indices, *J. Comput. Nonlinear Dyn.* 14 (11) (2019) 111008.